

De veneriske Sygdomme i Danmark udenfor Kjøbenhavn.

Tab. III.

	Gonorrhoe.	Venerisk Saar.	Syfilis.	Salt.
1864	1,483	533	704	2,720
1865	1,603	1,782		3,385
1866	1,604	478	1,167	3,249
1867	1,608	411	1,028	3,047
1868	1,667	516	966	3,149
1869	1,964	602	985	3,551
1870	1,936	405	748	3,089
1871	1,786	456	867	3,109
1872	1,925	663	856	3,444
1873	1,827	509	824	3,160
1874	1,760	564	734	3,058
1875	1,564	328	555	2,447
1864—1875	20,727	16,681		37,408
1876	1,770	367	405	2,542
1877	1,537	288	342	2,167
1878	1,472	224	341	2,037
1879	1,500	254	343	2,097
1880	1,623	244	352	2,219
1881	1,687	305	388	2,380
1882	1,655	383	461	2,499
1883	1,666	265	459	2,390
1884	1,615	255	400	2,270
1885	1,502	241	393	2,136
1886	1,344	228	373	1,945
1887	1,229	179	361	1,769
1876—87	18,600	3,233	4,618	26,451
		7,851		
1864—75 paa 10,000 af Befolkningen $\frac{1}{2}$ 1870 (1,603,450).	128	104		233
1876—87 paa 10,000 af en beregnet Befolkning d. $\frac{1}{2}$ 1882 (1,761,537).	105	45		150