

## Venertiske Sydomme i Kjøbenhavn.

Tab. IV.

Åar.	Beregnet Befolkning.	Gonorrhoe.	Veneriff Saar.	Syfilis.	Tilsammen.
1864	163,000	2,488	1,083	514	4,085
1865	166,000	2,656	836	577	4,069
1866	169,000	2,511	946	766	4,223
1867	173,000	2,248	886	869	4,003
1868	173,000	2,456	1,255	982	4,693
1869	176,000	2,631	1,639	1,019	5,289
1870	181,000	2,877	1,052	861	4,790
1871	190,000	2,737	898	770	4,405
1872	194,000	3,005	1,344	858	5,207
1873	197,000	2,817	1,366	743	4,926
1874	201,000	3,158	1,365	811	5,334
1875	211,000	3,335	837	690	4,862
1864—75		32,919	13,507	9,460	55,886
			22,967		
1876	213,000	3,843	1,154	696	5,693
1877	220,000	4,129	1,018	709	5,856
1878	224,000	4,426	845	717	5,988
1879	229,000	4,498	882	940	6,320
1880	233,000	4,538	968	954	6,460
1881	246,000	5,000	1,340	1,005	7,345
1882	252,000	5,654	1,533	1,077	8,264
1883	261,000	5,947	1,329	1,182	8,458
1884	269,000	5,673	1,670	1,341	8,684
1885	280,000	6,023	1,453	1,868	9,344
1886	290,000	5,844	1,590	2,122	9,556
1887	290,000	5,385	934	1,736	8,055
1876—87		60,960	14,716	14,347	90,023
			29,063		
1864—75 anmeldtes paa 10,000 af Befolk- ningen (181,000).		1,818	1,269		3,087
1876—87 anmeldtes paa 10,000 af Befolk- ningen (252,000).		2,419	1,153		3,572